



Possible complications after hand surgery

Reviewed: October 2024 Next review: October 2025

Version 4



The purpose of this leaflet is to provide information about the possible risks following hand surgery. With all procedures there are some risks. It is important that you understand the risks so that you can make a decision whether you want to have the procedure performed.



Swelling

Swelling is common in the operated area.

Elevating your hand and moving your fingers will help keep swelling down.

If the swelling increases and the affected area looks red, this may indicate an infection. If you suspect you are developing an infection, you should contact your General Practitioner or the Hand Clinic.

Bruising

Bruising is common after operations. There may be some minor seepage from your wound following the surgery and this will settle over time.

Bleeding

If you experience steady bleeding from your wound, elevate your hand and press on the wound to stop the bleeding. If it does not stop, depending on the severity of the bleed, contact your General Practitioner, Accident and Emergency or the Hand Clinic.

Infection

Superficial infections are common (5%) after hand surgery and cause redness, swelling, pain and pus around the stitches. These tend to settle quickly with oral antibiotics and regular dressings.

Deep infections around the tendons and bones are rare and may need admission to hospital for intravenous antibiotics and further surgery. These can cause longer-term complications, such as stiffness.

Scarring

Every operation leaves scars.

Your scar may initially be tender and thickened. It should soften and become more comfortable with time. Firm massage should help the scar improve.

Your scar may be numb permanently.

You will be provided you with information about hand care after surgery.

Delayed wound healing

Your wound will heal with dressings but may take a few weeks to heal completely.

Sometimes the wound splits open or takes longer to heal. The reasons for this may include infection, bleeding or poor blood supply to the wound edges.

Stiffness

Stiffness may be due to swelling, scar tissue or immobility. It can be made worse by infection or other complications. You can help to avoid stiffness by keeping your fingers moving after your surgery.

Damage to adjacent structures

Other nerves, vessels and structures could be inadvertently damaged during the operation. This is more common in operations that occur after trauma or when there is a lot of scar tissue at the site of the surgery.

Complex Regional Pain Syndrome (CRPS)

Complex Regional Pain Syndrome (CRPS) is chronic pain syndrome that causes pain, stiffness and swelling which are out of proportion to the surgery.

The pain may be experienced as burning and like an electric shock. If the pain is not controlled by simple painkillers, contact the Hand Clinic.

It is not fully understood and surgeons cannot predict who will be affected.

Mild CRPS occurs in 5% of patients and will have good resolution.

Severe cases are very rare but may leave the hand permanently stiff and swollen.

Cold intolerance

Cold intolerance is very common after procedures on your finger but the reason for this is not clear.

It develops over the first three months following surgery and may remain fairly constant over the first 12 months. It may improve over subsequent years but rarely resolves completely.

This may cause you difficulty working in cold environments. You can protect your hands from both wind-chill and cold by wearing insulated gloves.





If you have any questions or concerns, please email handsurgery@gjnh.scot.nhs.uk or contact the orthopaedic clinic during office hours on 0141 951 5521. For out of hours, you should contact NHS 24.



Post hand surgery videos

Scan the QR codes using your device or visit the web address listed below to view our videos on taking care of your hand after surgery. If you are viewing the digital version of this document, you can click the web address to go directly to the video.





All of our publications are available in different languages, larger print, braille (English only), audio tape or another format of your choice.

我們所有的印刷品均有不同語言版本、大字體版本、盲文(僅有英文)、錄音 帶版本或你想要的另外形式供選擇。

كافة مطبوعاتنا متاحة بلغات مختلفة و بالأحرف الطباعية الكبيرة و بطريقة بريل الخاصة بالمكفوفين (باللغة الإنكليزية فقط) و على شريط كاسيت سمعي أو بصيغة بديلة حسب خيارك.

Tha gach sgrìobhainn againn rim faotainn ann an diofar chànanan, clò nas motha, Braille (Beurla a-mhàin), teip claistinn no riochd eile a tha sibh airson a thaghadh.

हमारे सब प्रकाशन अनेक भाषाओं, बड़े अक्षरों की छपाई, ब्रेल (केवल अंग्रेज़ी), सुनने वाली कसेट या आपकी पसंदनुसार किसी अन्य फॉरमेट (आस्प) में भी उपलब्ध हैं। 我们所有的印刷品均有不同语言版本、大字体版本、盲文(仅有英文)、录音带版本或你想要的另外形式供选择。

ਸਾਡੇ ਸਾਰੇ ਪਰਚੇ ਅਤੇ ਕਿਤਾਬਚੇ ਵਗ਼ੈਰਾ ਵੱਖ ਵੱਖ ਭਾਸ਼ਾਵਾਂ ਵਿਚ, ਵੱਡੇ ਅੱਖਰਾਂ ਅਤੇ ਬ੍ਰੇਲ (ਸਿਰਫ਼ ਅੰਗਰੇਜ਼ੀ) ਵਿਚ, ਅੱਡੀਓ ਟੇਪ 'ਤੇ ਜਾਂ ਤੁਹਾਡੀ ਮਰਜ਼ੀ ਅਨੁਸਾਰ ਕਿਸੇ ਹੋਰ ਰੂਪ ਵਿਚ ਵੀ ਮਿਲ ਸਕਦੇ ਹਨ।

جاری تمام مطبوعات مختلف زبانوں، بوے حروف کی چیپائی، بریل (صرف اگریزی)، سنے والی کسٹ یا آپ کی پیند کے مطابق کسی دیگر صورت (فارمیٹ) میں بھی دستیاب ہیں۔



2: 0141 951 5513