



Reclining chair – supporting your recovery

Reviewed: March 2026
Next review: March 2027
Version 2

About this leaflet

The purpose of this leaflet is to introduce a new way to support your recovery.

Following your operation, you may be returned to a recliner chair. These have been proven to support your recovery in a number of ways including the following:

1. Shortened hospital stay

Patients that walk soon after surgery are discharged earlier than those who delay movement. This is because physical activity strengthens your muscles and joints, resulting in increased circulation which can accelerate your recovery.

2. Improved muscle strength

One of the many benefits of early mobilisation after surgery is improved muscle strength. Moving soon after your operation will help you regain muscle strength, which will help with your coordination, posture and balance. It will also aid in increased joint flexibility in your knees, hips and ankles. Each of these benefits will get you closer to regaining independent mobility soon after your operation.

3. Increased blood flow

Early mobilisation after surgery increases blood flow. Increased blood flow offers several benefits that minimise the risk of complications and help your body heal faster.

Increased blood flow stimulates circulation and reduces the risk of blood clots. Post-operative patients are naturally at a high risk for blood clots due to decreased blood flow.

This also aids in quicker wound healing, reduced swelling and helps maximise nutrients and oxygen delivered to your surgical site. Each of these factors will help accelerate the healing process.

4. Improved independence

Early mobilisation after surgery will accelerate your return to independent mobility as you patiently regain your strength. Improved independence is known to improve patients' moods and self-esteem, further aiding in a positive healing journey.

5. Improved physical function

Early mobilisation after surgery will also improve your physical function. Many of your body systems are naturally slowed after surgery.

6. Less post-operative complications

Remember, without early mobilisation, patients are at a higher risk for postoperative complications. Staying in bed for an extended period of time after surgery leads to issues such as:

- Mobility and functional restrictions
- Sleep deprivation
- Delirium
- Altered nutritional states
- Prolonged hospitalisation

7. Improved patient outcomes

Early mobilisation has been proven to enhance patient outcomes. It minimises complications associated with bedrest, improves bodily functions, increases strength and endurance, and encourages a positive outlook.

➤ Contact

If you have any questions about your recover please speak to a member of the nursing team.

All of our publications are available in different languages, larger print, braille (English only), audio tape or another format of your choice.

我們所有的印刷品均有不同語言版本、大字體版本、盲文（僅有英文）、錄音帶版本或你想要的另外形式供選擇。

كافة مطبوعاتنا متاحة بلغات مختلفة و بالأحرف الطباعية الكبيرة و بطريقة بريل الخاصة بالمكفوفين (باللغة الإنكليزية فقط) و على شريط كاسيت سمعي أو بصيغة بديلة حسب خيارك.

Tha gach sgrìobhainn againn rim faotainn ann an diofar chànanan, clò nas motha, Braille (Beurla a-mhàin), teip clàistinn no riochd eile a tha sibh airson a thaghadh.

हमारे सब प्रकाशन अनेक भाषाओं, बड़े अक्षरों की छपाई, ब्रेल (केवल अंग्रेज़ी), सुनने वाली कसेट या आपकी पसंदनुसार किसी अन्य फॉरमेट (आरएफ) में भी उपलब्ध हैं।

我們所有的印刷品均有不同語言版本、大字體版本、盲文（僅有英文）、錄音帶版本或你想要的另外形式供選擇。

ਸਾਡੇ ਸਾਰੇ ਪਰਚੇ ਅਤੇ ਕਿਤਾਬਚੇ ਵਗੈਰਾ ਵੱਖ ਵੱਖ ਭਾਸ਼ਾਵਾਂ ਵਿਚ, ਵੱਡੇ ਅੱਖਰਾਂ ਅਤੇ ਬ੍ਰੇਲ (ਸਿਰਫ਼ ਅੰਗਰੇਜ਼ੀ) ਵਿਚ, ਆਡੀਓ ਟੇਪ 'ਤੇ ਜਾਂ ਤੁਹਾਡੀ ਮਰਜ਼ੀ ਅਨੁਸਾਰ ਕਿਸੇ ਹੋਰ ਰੂਪ ਵਿਚ ਵੀ ਮਿਲ ਸਕਦੇ ਹਨ।

ہماری تمام مطبوعات مختلف زبانوں، بڑے حروف کی چھپائی، بریل (صرف انگریزی)، سنے والی کست یا آپ کی پسند کے مطابق کسی دیگر صورت (فارمیٹ) میں بھی دستیاب ہیں۔



: 0141 951 5513