



Trigger finger and trigger thumb

Reviewed: October 2024 Next review: October 2025

Version 4

About this leaflet

The purpose of this leaflet is to tell you about Trigger Finger and Trigger Thumb and how they can be treated.

About Trigger Finger and Trigger Thumb

Trigger Finger is a condition where a finger or thumb clicks, catches or locks as it bends and straightens.

Symptoms

- Pain in the palm of the hand.
- Tender when pressed.
- Finger clicks with movement or gets stuck in a bent position.
- Stiffness which causes difficulty bending the thumb.
- Finger becomes stuck in a bent position and you are unable to release it yourself. If this happens, you will need to seek urgent medical attention so the joint does not become permanently stiff.

Causes

The tendons that bend your fingers run into a tunnel. If the mouth of the tunnel becomes thickened, the tendon can swell, causing it to catch in the mouth of the tunnel. People with diabetes are more prone to getting trigger finger.

Diagnosing Trigger Finger and Trigger Thumb

Diagnosis is usually made by:

- · talking about your symptoms and the troubles you are having; and
- examining your hand.

Treatment options

Trigger Finger does not usually cause harm, however it can be a nuisance and painful.



Non-surgical treatment

Some mild cases resolve without treatment. There are a number of non-surgical treatment options.

- Splint: Putting a splint on the finger to keep it straight at night keeps the swollen area of the tendon inside the tunnel and help it smooth back down. You could use a simple lollipop stick as a home-made splint.
- Steroid injection: A steroid injection into the affected finger or thumb can improve your symptoms. This is effective in about 70% of people, but the success rate is lower if you have diabetes. The injection can be carried out in the outpatient clinic. You would be able to drive after an injection.

If your symptoms persist despite non-surgical treatment, surgery may be an option.

Surgical treatment – Trigger Finger or Trigger Thumb Release Surgery

The aim of the operation is to open the entrance to the tunnel that your finger tendon runs on to let it glide freely. The operation takes about 10 minutes to perform so is carried out under local anaesthetic as a day case.

The palm of your hand near the finger/thumb will be numbed with local anaesthetic. You should not feel pain during the operation but may feel some pressure and your finger/thumb might go numb.

A tight band called a tourniquet is placed on your upper arm to stop the blood flow for a few minutes so that the surgeon has a good view to perform the operation.

The surgeon will cut through the skin, divide the entrance of the tunnel, make sure that there is no more catching of the finger and then stitch you up.

Risks

- Tender scar: This can last for a few months and can be improved by massaging the scar. You will be shown how to do this before you go home.
- Swelling: Your finger will feel a bit swollen and stiff for a while after the surgery.
- Infection and Bleeding: There is a small risk of infection or bleeding.
- Recurrence: Although it is uncommon to get triggering again in the same finger/thumb, some people develop it in other fingers/thumbs.
- Numbness: The risk of a serious severe complication is very low (<1%). The nerves that supply
 your finger are very close to the tendon and there is a very small chance that a nerve could be
 injured. This could leave you with some numbness in the finger or a painful nerve swelling called
 a neuroma.
- Complex Regional Pain Syndrome: This is a very rare condition that can occur after any hand surgery. It causes stiffness, swelling and pain out of proportion to the procedure. It is like a reflex that hasn't switched off. This can be treated with pain relief and physiotherapy.

Recovery period

You will have stitches in for 10-14 days. It is really important that you keep your fingers moving throughout this time. You can use your hand for light tasks during this period.

You should check with your insurance company before drive with stitches in your hand.



If you decide not to have treatment

Sometimes your symptoms may resolve with time.

If your finger becomes stuck in a bent position and you are unable to straighten it yourself you should seek medical attention so that the joint does not become permanently stiff.



Contact

If you have any questions or concerns, please email handsurgery@gjnh.scot.nhs.uk or contact the orthopaedic clinic during office hours on 0141 951 5521. For out of hours, you should contact NHS 24.



Post hand surgery videos

Scan the QR codes using your device or visit the web address listed below to view our videos on taking care of your hand after surgery. If you are viewing the digital version of this document, you can click the web address to go directly to the video.





All of our publications are available in different languages, larger print, braille (English only), audio tape or another format of your choice.

我們所有的印刷品均有不同語言版本、大字體版本、盲文(僅有英文)、錄音 帶版本或你想要的另外形式供選擇。

كافة مطبو عاتبًا متاحة بلغات مختلفة و بالأحرف الطباعية الكبيرة و بطريقة بريل الخاصة بالمكفوفين (باللغة الإنكليزية فقط) و على شريط كاسيت سمعى أو بصيغة بديلة حسب خيارك.

Tha gach sgrìobhainn againn rim faotainn ann an diofar chànanan, clò nas motha, Braille (Beurla a-mhàin), teip claistinn no riochd eile a tha sibh airson a thaghadh.

हमारे सब प्रकाशन अनेक भाषाओं, बड़े अक्षरों की छपाई, ब्रेल (केवल अंग्रेज़ी), सुनने वाली कसेट या आपकी पसंदनुसार किसी अन्य फॉरमेट (आस्प) में भी उपलब्ध हैं। 我们所有的印刷品均有不同语言版本、大字体版本、盲文(仅有英文)、录音带版本或你想要的另外形式供选择。

ਸਾਡੇ ਸਾਰੇ ਪਰਚੇ ਅਤੇ ਕਿਤਾਬਚੇ ਵਗ਼ੈਰਾ ਵੱਖ ਵੱਖ ਭਾਸ਼ਾਵਾਂ ਵਿਚ, ਵੱਡੇ ਅੱਖਰਾਂ ਅਤੇ ਬ੍ਰੇਲ (ਸਿਰਫ਼ ਅੰਗਰੇਜ਼ੀ) ਵਿਚ, ਆੱਡੀਓ ਟੇਪ 'ਤੇ ਜਾਂ ਤੁਹਾਡੀ ਮਰਜ਼ੀ ਅਨੁਸਾਰ ਕਿਸੇ ਹੋਰ ਰੂਪ ਵਿਚ ਵੀ ਮਿਲ ਸਕਦੇ ਹਨ।

ہاری تمام مطبوعات مختلف زبانوں، بو ہے حروف کی چھپائی، بریل (صرف انگریزی)، سنے والی کسٹ یا آپ کی پیند کے مطابق کسی دیگر صورت (فارمیٹ) میں بھی دستیاب ہیں۔

