



Taking opioid pain relief following surgery

Reviewed: May 2024
Next review: May 2025
Version 1

> About this booklet

This booklet tells you about **opioids** used for pain relief in useful sections.

> About opioid pain relief

Opioids provide pain relief by acting on areas in the spinal cord and brain to block the transmission of pain signals. They are considered to be some of the strongest painkillers available and are used to treat pain after surgery, serious injury and cancer. Strong opioids include Morphine, Oxycodone and Fentanyl.

Only take them if other pain medicines are unsuitable or are not providing enough pain relief.

Prescribed in either slow release or fast release form.

If you were taking strong opioid medication before coming into hospital, you may have been advised to temporarily increase your usual dose after your surgery.

Only use opioids short term.

Information is available in your patient information booklet.

Dosage of any medicine is the lowest dose that produces the desired benefit.

Side effects from strong opioids

Only take them if other pain medicines are unsuitable or are not providing enough pain relief.

You are unlikely to be entirely pain free after surgery. The aim of pain management after surgery is to provide enough pain relief to allow you to function reasonably comfortably and keep side-effects to a minimum. You will be given a combination of several different types of painkillers, depending on what is most suitable for you.

Prescribed in either slow release or fast release form.

Slow release tablets

The painkiller in slow release opioids (tablets or capsules) is released slowly into your body over 12 hours.

Fast release tablets

The fast release opioid form (tablet or liquid) will have a noticeable effect within approximately 30 minutes and last up to four hours.

If you were taking strong opioid medication before coming into hospital, you may have been advised to temporarily increase your usual dose after your surgery. In this instance, it is important to try to reduce this to your usual dose in the time frame advised when you were discharged, or sooner if you feel able.

If your surgery has fixed the cause of your pain, you should speak to your General Practitioner (GP) about how to slowly reduce and stop your long term opioids to avoid experiencing withdrawal symptoms.

Only use opioids short term. Your GP will supervise a gradual reduction to help you stop taking them as your surgical pain improves.

The risk to your health increases significantly when using strong opioids for a long period of time.

You can become dependent on them. This means that if you stop taking them suddenly, you can get withdrawal symptoms such as tiredness, sweating, runny nose, stomach cramps, diarrhoea and aching muscles.

Information is available in your patient information booklet. If you have any questions about pain relief before you go home, you can ask the ward nurses looking after you, ward pharmacist or the specialist pain nurses.

If you have any questions after you go home, contact the follow up service number in your surgery specific information booklet.



Dosage of any medicine is the lowest dose that produces the desired benefit.

It is not usual to get complete relief of pain from opioids. The aim of treatment is to reduce your pain enough for you to be able function reasonably comfortably as you recover.

We will give you advice on the suitable dose for you.

When to take opioid medicines:

Slow release tablets

These are designed to produce a steady level of opioid medication in your blood over 12 hours. You should take them twice a day, 12 hours apart, for example at 6am and 6pm.

It is important to swallow the tablet or capsule whole – do not chew or crush them.

If you miss a dose of slow release opioid take it as soon as you remember. However, if it is almost time for your next dose, skip the missed dose and take your medication as normal. Do not take two doses at the same time.

Fast release tablets

You can take fast release opioid tablets four to six hours apart but with no more than four doses in 24 hours.

You can use them to top-up your pain relief if needed in between taking your other pain relief medication.

We will advise you how long to take your strong opioids for. You should discuss with your GP before your supply runs out.

Side effects from strong opioids

Side effects that can be experienced when taking opioid medication include:

- Dizziness
- Constipation
- Nausea (feeling sick)
- Vomiting (being sick)
- Feeling sleepy
- Confusion
- Itching
- Difficulty breathing at night

Your healthcare team may prescribe:

- anti-sickness medication to help reduce nausea laxatives to help treat constipation.

Side effects that are more likely to occur if you need to continue taking opioid medication beyond the first few days after your surgery may include:

- addiction
- weight gain
- erectile dysfunction
- lack of sex drive
- depression
- reduced immunity
- irregular periods
- increased levels of pain
- tolerance (they become less effective at relieving pain)

Driving

It is an offence to drive while impaired by medication.

You are responsible for making sure your ability to drive is not impaired and that you are safe to drive each time you do so. Your surgery itself, pain and the side effects of opioids can all impair your ability to drive.

It is not advised to drive within 24 hours of a general anaesthetic.

Alcohol

Drinking alcohol while taking opioids can cause sleepiness and poor concentration. You should avoid alcohol completely when you first start taking opioids or if you have recently increased your usual dose.

Discarding unused medication

Opioids can be harmful if taken by adults, children or animals.

Unused tablets should be taken to your local pharmacy for disposal.

Do not put unused medicines in household waste flush them down the toilet or put them down the sink.

Further information

If you have any questions after you go home, contact the follow up service number in your surgery specific information booklet.

All of our publications are available in different languages, larger print, braille (English only), audio tape or another format of your choice.

我們所有的印刷品均有不同語言版本、大字體版本、盲文（僅有英文）、錄音帶版本或你想要的另外形式供選擇。

كافة مطبوعاتنا متاحة بلغات مختلفة و بالأحرف الطباعية الكبيرة و بطريقة بريل الخاصة بالمكفوفين (باللغة الإنكليزية فقط) و على شريط كاسيت سمعي أو بصيغة بديلة حسب خيارك.

Tha gach sgrìobhainn againn rim faotainn ann an diofar chànanan, clò nas motha, Braille (Beurla a-mhàin), teip clàistinn no riochd eile a tha sibh airson a thaghadh.

हमारे सब प्रकाशन अनेक भाषाओं, बड़े अक्षरों की छपाई, ब्रेल (केवल अंग्रेज़ी), सुनने वाली कसेट या आपकी पसंदनुसार किसी अन्य फॉरमेट (आरप) में भी उपलब्ध हैं।

我們所有的印刷品均有不同語言版本、大字體版本、盲文（僅有英文）、錄音帶版本或你想要的另外形式供選擇。

ਸਾਡੇ ਸਾਰੇ ਪਰਚੇ ਅਤੇ ਕਿਤਾਬਚੇ ਵਗੈਰਾ ਵੱਖ ਵੱਖ ਭਾਸ਼ਾਵਾਂ ਵਿਚ, ਵੱਡੇ ਅੱਖਰਾਂ ਅਤੇ ਬ੍ਰੇਲ (ਸਿਰਫ਼ ਅੰਗਰੇਜ਼ੀ) ਵਿਚ, ਆਡੀਓ ਟੇਪ 'ਤੇ ਜਾਂ ਤੁਹਾਡੀ ਮਰਜ਼ੀ ਅਨੁਸਾਰ ਕਿਸੇ ਹੋਰ ਰੂਪ ਵਿਚ ਵੀ ਮਿਲ ਸਕਦੇ ਹਨ।

ہماری تمام مطبوعات مختلف زبانوں، بڑے حروف کی چھپائی، بریل (صرف انگریزی)، سنے والی کست یا آپ کی پسند کے مطابق کسی دیگر صورت (فارمیٹ) میں بھی دستیاب ہیں۔



: **0141 951 5513**