



Thoracic prehabilitation

– prepare for your lung surgery or treatment

Reviewed: April 2026
Next review: April 2027
Version 5

➤ About this leaflet


The purpose of this leaflet is to provide patients with information about safe levels of activity and how to improve their health and levels of fitness before treatment or surgery.

You may have been given a date for surgery or are waiting to hear if surgery is required. The information is applicable in both of these cases.

➤ What is prehabilitation and why is it important?

Prehabilitation is designed to maximise an individual's overall general health and level of fitness to enable them to withstand the effects of treatment or surgery and introduces steps you can take to help you prepare for the next step in your treatment journey. The actions you take now can help you recover quicker, reduce the risk of complications and reduce the time you spend in hospital. Many people are happy to know they can do something to improve their health in the lead up to surgery or treatment.

You may only have a short time between potential diagnosis and your treatment/surgery, however, small changes can make a big difference. The benefits of prehabilitation can be seen in as little as 7-10 days.




Prehabilitation means your journey to recovery starts before your treatment/surgery has even begun through physical, nutritional and psychological support.

➤ **Benefits of prehabilitation**

- Reduce length of stay in hospital
- Reduce complications after surgery
- Improve physical function
- Improve overall fitness
- Improve emotional wellbeing
- Improve nutritional health
- Provide a sense of control and purpose
- Enhance quality of life
- Promote long-term positive health behaviour

Prehabilitation is not a standalone treatment, it consists of many key components to help you 'Get READY' for surgery:

Get **READY**



Respiratory exercises



Eat well



Activity



Dental and oral hygiene



Your lifestyle and wellbeing

▶ Respiratory exercises

Below are the step by step instructions for the deep breathing exercise. Aim to do this exercise 3 times a day, ideally morning, noon and in the evening. These are the same exercises you will carry out post operatively with the physiotherapist.

Positioning is important: before you start, make sure you are sitting upright and not in a slumped position. Take a couple of practice breaths with your hands on your stomach – you should feel your hands being pushed out as you inhale and move in as you exhale.

Deep breathing exercise



These are the step-by-step instructions for your deep breathing exercise. If you can, carry out this exercise in a high sitting position.



◀ Step 1

Take a deep breath in through your nose, hold for 3 seconds and breath out slowly through your mouth.

Repeat 3 times.

Step 2 ▶

Huff – sharply breathe out through your mouth. This is the same action as using your breath to steam up a mirror.

Repeat 3 times.



◀ Step 3

Cough.

Scan the QR code to view our breathing exercises video



Incentive Spirometer and Inspiratory Muscle Trainer

Below are step by step instructions for using your Incentive Spirometer and Inspiratory Muscle Trainer.

It is recommended you use your Incentive Spirometer 2 to 3 times per day to record a measurement of your inspiratory breath.

Chest training: Incentive Spirometer

- Step 1: Sit upright
- Step 2: Hold the device in an upright position
- Step 3: Put the mouthpiece in your mouth and close your lips around it
- Step 4: Breathe in slowly through your mouth for as long as you can. The green piston should rise the middle column. The green button should stay between the arrows.
- Step 5: Keep the green button floating for as long as you can and then breathe out slowly.
- Step 6: Rest for a few seconds and repeat steps 1-5 for 3-5 breaths. Record your highest breath daily in the diary provided.



Scan the QR code to view the incentive spirometer video

Please bring spirometer in to hospital as you can continue to use it after surgery.



Inspiratory Muscle Training (POWER®breathe)

Inspiratory Muscle Training (IMT) is a form of resistance training for the muscles that we use to breathe.

The resistance from the device makes the intercostal muscles (between the ribs) and the diaphragm (under the lungs) work harder for short periods of time. This makes them stronger and can make breathing easier.

Doing IMT before an operation can reduce the risk of breathing complications after surgery.

1. Make sure you are in an upright sitting position or standing.
2. Hold the hand grip and put the mouthpiece in your mouth, sealing it with your lips.
3. Breathe out if you can then take a fast, hard breath in through your mouth. Try and take as much air into your lungs as possible.
4. Breathe out slowly, relaxing your shoulders.
5. Take 2 relaxed breaths to recover and repeat steps 3 and 4.
6. If you feel lightheaded, stop and take time to settle before continuing. If this does not settle, stop using the device and seek medical advice.

The aim is to complete 2 sets of 10 deep breaths with the device twice a day and build up the resistance level over time. You can rest as much as you need until you become comfortable using the device.


It may take you some time to build up and you should only increase the resistance if you have built up to 2 sets of 10 twice per day and are finding this easy.

Record the level of IMT in the diary provided so you can see your progress.

Caring for the device: Cleaning advice and sterile solution is included with the device.

Scan the QR code to view the inspiratory muscle trainer video

Please bring IMT with you to hospital as you can continue to use after surgery.



Eating a balanced diet is an important part of maintaining good health to help prepare for your surgery. The Eatwell Guide shows you how much of what you eat should come from each food group.

Fruit and vegetables: Aim to eat at least 5 portions of a variety of fruit and vegetables each day.

Bread, rice, pasta or other starchy carbohydrates: Starchy food should make up just over a third of the food we eat. Choose higher fibre wholegrain varieties, such as wholewheat pasta and brown rice, or simply leave skins on potatoes.

Milk and dairy products: Good sources of protein and vitamins. They are an important source of calcium too. Try to choose low fat varieties to limit your intake of saturated fat.

Meat, fish, eggs and beans: These foods are good sources of protein, vitamins and minerals. Pulses, such as beans, peas and lentils are good alternatives to meat because they are lower in fat and higher in fibre and protein too. Choose lean, low fat, cuts of meat and mince, and eat less red and processed meat like bacon, ham and sausages. Aim for at least 2 portions of fish every week, 1 of which should be oily, such as salmon or mackerel.


Choose unsaturated oils and spreads, and eat in small amounts: Unsaturated fats are healthier fats and include vegetable, rapeseed, olive and sunflower oils. Remember all types of fat are high in energy, but should be eaten sparingly.

Information source: <https://www.nhs.uk/live-well/eat-well/the-eatwell-guide/>

For people who are underweight, have recently lost weight unintentionally or have a poor appetite, it is important to maintain a stable weight and prevent further weight loss prior to your surgery.

The following ideas may help you to increase your energy and protein intake:

- Eat little and often. Aim to have 3 meals and 3 snacks per day
- Avoid 'low fat', 'sugar-free', or 'diet' products
- Choose the meals you enjoy
- Have a nourishing dessert/pudding – full fat yogurt, rice pudding or sponge and custard
- Have a pint of full fat milk everyday – for example on cereal, milky drinks or in puddings
- Add butter or margarine, cream or cheese to mashed potatoes, vegetables, scrambled eggs and hot pasta
- Have drinks (tea/coffee/juice) with or after meals as these can fill you up if taken just before food.



There is strong evidence that it is safe to stay active in the time leading up to surgery or treatment. You should try and do 30 minutes of moderate level activity every day. If you cannot manage this in one go, try breaking the activity into smaller manageable periods, for example, 10-15 minutes activity in the morning and then 10-15 minutes in the afternoon.

Physical activity does not mean a lot of hard work or expense, it could be going out for a walk, gardening, or using your stairs at home. Avoid sitting for long periods of time. Your physiotherapist will issue you with a personalised exercise programme at your appointment. You may notice changes to your breathing when exercising. This can show if you are exercising to an appropriate level.

You should aim to be a little breathless, but still be able to hold a conversation with someone. If you are too breathless, to do this you must stop, catch your breath and then carry on. You should aim to be working to level 3-5 on the Rate of Perceived Exertion (RPE) scale in this booklet.

I already exercise and am quite fit. What should I do?

If you already exercise regularly and your Consultant is happy for you to do so, then please continue with your routine. Keeping up your fitness will help you cope with the effects of your surgery better and exercise may also help you to cope with the worry and anxiety upcoming surgery brings. Please discuss with a healthcare professional involved in your care if you have any concerns regarding activities or exercises.

Walking

Walking is often overlooked as a form of exercise, but it is simple, free and one of the easiest ways to get more active and healthier. A brisk 10-minute daily walk has a range of health benefits and counts towards your recommended 150 minutes of weekly exercise.

You can tell you are walking briskly if you can talk but cannot sing. If you become breathless enough that you cannot talk to someone, stop, catch your breath and carry on. Your physiotherapist will provide you with a Rate of Perceived Exertion (RPE) scale to guide you.

Tips:

- Plan a route in advance.
- Remember wherever you are going, you must be able to get back.
- Take a mobile phone or inform someone you are going for a walk.
- Get others involved!
- Use a walking aid if necessary.
- Wear comfortable, supportive shoes.

You can monitor your daily step count using a pedometer. Most mobile phones have a pedometer or step count function included. It is recommended that you increase your step count by 10% each week. Please fill in the step count diary to monitor your progress.

Rated Perceived Exertion (RPE) Scale

0 At rest (no exertion at all)

0.5 Very easy (very, very light)

1 Easy (very light)

2 Not difficult (fairly light)

3

4 A little hard

5

6 Hard

7

8 Very hard

9

10 Extremely hard



Main exercises:


Warm up for 1 minute marching on the spot in standing or in sitting

Sit to stand:



Sitting in a chair and keeping your feet hip width apart, push on the arms of the chair to stand up. Slowly and controlled, sit back down. Repeat 10 times.

Heel raises:



Standing freely or holding on to a wall/chair for support, rise up to your toes. Hold here for 3 seconds and slowly release your heels back down to the floor. Repeat 10 times.

Mini squats:




Standing freely or using a chair/wall for support, slowly bend your knees to a comfortable angle and release back to standing. Repeat 10 times.

Hip abductions:



Standing freely or using a chair/wall for support, lift your leg out to the side, keeping your knee straight. Repeat 10 times with each leg.

Hip extensions:



Standing freely or using a chair/wall for support, lift your leg out behind you, keeping your knee straight. Repeat 10 times with each leg.

Shoulder raises:



Take a deep breath in and on exhale slowly raise one or both arms straight up in line with your ears. Slowly drop them back down by your side. Repeat 10 times.

Cool down:

It is very important to complete a cool down after exercise to help restore your body to a resting state. After completing your exercise programme, you should march on the spot for around 3-5 minutes, gradually slowing your pace.

In addition, complete a cycle of the deep breathing exercise on page 3.

Daily diary

Please record the number of steps you have achieved for each day using a pedometer/ phone/ smart watch, record the level your inspiratory muscle trainer (IMT) is set at and record the highest breath achieved on your incentive spirometer.

Week 1	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 2	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 3	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 4	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 5	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							



Week 6	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 7	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 8	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 9	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 10	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 11	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							
Week 12	Day	M	T	W	T	F	S	S
	Highest spirometer							
	Daily step count							
	IMT level							



Dental and oral hygiene

Good dental and oral hygiene reduces the risk of lung complications after surgery and improves your overall general health. You should aim to:

- Brush your teeth for two minutes, at least twice a day
- Use a fluoride mouthwash
- Use dental floss

For more information on recommendations for dental and oral hygiene including teeth cleaning guide, please refer to the NHS Inform website: www.nhsinform.scot

Your lifestyle and wellbeing

As well as eating well, it is also important to consider your lifestyle in relation to smoking and alcohol.

Stopping smoking/vaping

Giving up smoking/vaping is one of the most important steps in getting healthy before your operation. The sooner you stop smoking/vaping prior to surgery the better.

Smoking/vaping significantly increases risks in thoracic surgery by causing pulmonary complications (pneumonia, respiratory failure), cardiovascular events, and impaired wound healing. Active smokers face almost double the complication rate compared to non-smokers.

Quitting at least 4–8 weeks prior is strongly advised to mitigate these risks but quitting even a few days before surgery can reduce carbon monoxide levels, though longer is better.

To get help stopping smoking/vaping:








- Speak to your local pharmacist
- Call Quit Your Way 0800 84 84 84



Alcohol

You should also limit the amount of alcohol you drink. Do not exceed national recommendations, which are currently 14 units per week for both men and women. People who drink alcohol above the recommended amount are 3 to 5 times more likely to develop infections. They are also more likely to spend a longer time in intensive care, have problems with bleeding and have irregular heart rhythms after surgery.

Reducing alcohol intake one month before surgery can significantly reduce these risks. Reducing intake less than one month before can still be of benefit but to a lesser extent.

						
2.2 units	1.5 units		2.2 units	3.1 units	9.8 units	1 unit
1 pint of beer/lager/ cider (4% ABV)	330ml bottle of beer/lager/ cider (4.5% ABV)		175ml glass of wine (12.5% ABV)	250ml glass of wine (12.5% ABV)	750ml bottle of wine (13% ABV)	25ml measure of spirits (40% ABV)

For more information on healthy lifestyle choices, visit: www.nhsinform.scot

What does the programme involve?

The Physiotherapist will complete an initial assessment where you will be asked some questions and to complete some exercise tests to establish your current level of fitness. They will create a home exercise programme based on your initial assessment which will be made up of cardiovascular, strengthening and lung training exercises as appropriate.

We hope you find Prehabilitation gives you the guidance you can use to positively impact the time leading up to your surgery and help you take control of your health and wellbeing.

Post op advice and information

The physiotherapist will see you the day after the operation and you will be out of bed sitting in a chair and start walking under the supervision of the nurses and physiotherapists.

You should start your breathing exercises as soon as possible after surgery to help clear any phlegm from your chest and should advise your physiotherapist if you have an incentive spirometer or IMT device as you can restart this after surgery.

Please ensure that your nightwear has buttons if possible and that you have suitable footwear with supportive backs as you will be up walking the day after surgery.

Further information

Your healthcare team are here to support you. Any question that is important to you is important to us. Please contact the prehabilitation physiotherapist on **0141 951 5121** or email gjnh.heartandlungprehab@gjnh.scot.nhs.uk if you need help or advice to discuss any of the information in this booklet.

Thoracic Surgery Patient Journey - Scan the QR code or follow the link to watch a short video about the Thoracic Surgery Patient Journey, explaining what to expect before, during, and after your surgery.

<https://youtu.be/ng6WOa6nGjE>



NHS Scotland Prehabilitation for Scotland - resources to help you prepare for surgery.

<https://www.prehab.nhs.scot/prehab-and-me/>



Maggie's Centre - Maggie's Centre website offers free practical, emotional, and social support for people affected by cancer, including patients, families, and carers.

<https://www.maggies.org/>



All of our publications are available in different languages, larger print, braille (English only), audio tape or another format of your choice.

我們所有的印刷品均有不同語言版本、大字體版本、盲文（僅有英文）、錄音帶版本或你想要的另外形式供選擇。

كافة مطبوعاتنا متاحة بلغات مختلفة و بالأحرف الطباعية الكبيرة و بطريقة بريل الخاصة بالمكفوفين (باللغة الإنكليزية فقط) و على شريط كاسيت سمعي أو بصيغة بديلة حسب خيارك.

Tha gach sgrìobhainn againn rim faotainn ann an diofar chànanan, clò nas motha, Braille (Beurla a-mhàin), teip clàistinn no riochd eile a tha sibh airson a thaghadh.

हमारे सब प्रकाशन अनेक भाषाओं, बड़े अक्षरों की छपाई, ब्रेल (केवल अंग्रेज़ी), सुनने वाली कसेट या आपकी पसंदनुसार किसी अन्य फॉरमेट (आरएफ) में भी उपलब्ध हैं।

我們所有的印刷品均有不同語言版本、大字體版本、盲文（僅有英文）、錄音帶版本或你想要的另外形式供選擇。

ਸਾਡੇ ਸਾਰੇ ਪਰਚੇ ਅਤੇ ਵਿਤਾਬਚੇ ਵਗੈਰਾ ਵੱਖ ਵੱਖ ਭਾਸ਼ਾਵਾਂ ਵਿਚ, ਵੱਡੇ ਅੱਖਰਾਂ ਅਤੇ ਬ੍ਰੇਲ (ਸਿਰਫ਼ ਅੰਗਰੇਜ਼ੀ) ਵਿਚ, ਆਡੀਓ ਟੇਪ 'ਤੇ ਜਾਂ ਤੁਹਾਡੀ ਮਰਜ਼ੀ ਅਨੁਸਾਰ ਕਿਸੇ ਹੋਰ ਰੂਪ ਵਿਚ ਵੀ ਮਿਲ ਸਕਦੇ ਹਨ।

ہماری تمام مطبوعات مختلف زبانوں، بڑے حروف کی چھپائی، بریل (صرف انگریزی)، سنے والی کست یا آپ کی پسند کے مطابق کسی دیگر صورت (فارمیٹ) میں بھی دستیاب ہیں۔



 : **0141 951 5513**