



Post operative instructions following minor surgery

Reviewed: August 2026
Next review: August 2027
Version 9

About this leaflet

The purpose of this leaflet is to give you information following your surgery.

After surgery

Rest this evening and avoid strenuous activities. If you have had a general anaesthetic, in the next 24 hours you must not:

- drive;
- drink alcohol;
- use non-prescription drugs (other than simple painkillers); or
- use machinery.

If you have had local anaesthetic injections, it may take several hours for the numbness to wear off. Take your usual medications and any that you may have been given by the nurse today.

It is normal to feel some pain after surgery. You may be prescribed painkillers, otherwise take something simple, e.g. Paracetamol if required. You will be further advised before discharge.

You may feel drowsy or weak for several hours after your surgery. It may take 24 hours for the effects of a general anaesthetic to wear off completely.

A competent adult should stay with you overnight. You should eat and drink normally as able (unless otherwise advised).

Common side effects are:

- dizziness;
- drowsiness;
- nausea/vomiting;
- sore throat; and
- general aches and pains.

Investigations

If you have had a skin biopsy / or other investigations carried out, it can take between 4 to 6 weeks for the results of the specimen to be available to your GP practice.

Whilst your GP and ourselves try to ensure that any necessary action is taken when these results are available, you should personally check the outcome. Please contact your GP in the first instance. If there are any difficulties please contact the Golden Jubilee University National Hospital.

Wound care

Observe wound site for:

- bleeding or unexpected drainage that does not stop; and
- extreme redness or swelling around wound or drainage of pus.

Your sutures and/or dressings may have to be removed by your:

- Doctor's surgery; or
- Outpatient clinic;

You will be advised of this at discharge.

You may have been advised by the surgeon or a physiotherapist to:

- elevate affected arm or leg when resting and overnight;
- do specific exercises; and
- support wound when moving around or when coughing.

Please follow their instructions at home.

Other advice

You should be advised of a follow up appointment before you leave the hospital.

Contact the hospital if:

- you have severe pain not relieved by medication;
- you have extreme redness or swelling around the wound site or drainage of pus;
- you are unable to pass urine;
- you have continual vomiting; or
- you have a fever.

If your wound starts to bleed once you have been discharged home, you can apply pressure to the site for ten minutes. Keep the pressure applied with something that is dry and clean like a tissue, clean cloth, handkerchief or swabs.

If it does not stop bleeding after ten minutes, apply a cold compress, e.g. a bag of frozen peas wrapped in a pillowcase or towel for 20 minutes. If the bleeding does not stop or is excessive, contact your GP or NHS 24 for further advice.



This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.



All of our publications are available in different languages, larger print, braille (English only), audio tape or another format of your choice.

كافة مطبوعاتها متاحة بلغات مختلفة و بالأحرف الطباعية الكبيرة و بطريقة بريل الخاصة بالمكفوفين (باللغة الإنكليزية فقط) و على شريط كاسيت سمعي أو بصيغة بديلة حسب خيارك.

हमारे सब प्रकाशन अनेक भाषाओं, बड़े अक्षरों की छपाई, ब्रेल (केवल अंग्रेज़ी), सुनने वाली कसेट या आपकी पसंदनुसार किसी अन्य फॉरमेट (आरूप) में भी उपलब्ध हैं।

ਸਾਡੇ ਸਾਰੇ ਪਰਚੇ ਅਤੇ ਕਿਤਾਬਚੇ ਵਗ਼ੈਰਾ ਵੱਖ ਵੱਖ ਭਾਸ਼ਾਵਾਂ ਵਿਚ, ਵੱਡੇ ਅੱਖਰਾਂ ਅਤੇ ਬ੍ਰੇਲ (ਸਿਰਫ਼ ਅੰਗਰੇਜ਼ੀ) ਵਿਚ, ਅੱਡੀਓ ਟੇਪ 'ਤੇ ਜਾਂ ਤਹਾਫ਼ੀ ਮਰਜ਼ੀ ਅਨੁਸਾਰ ਕਿਸੇ ਹੋਰ ਰੂਪ ਵਿਚ ਵੀ ਮਿਲ ਸਕਦੇ ਹਨ।



☎: 0141 951 5513